DIFFUSIVE LIMIT OF ASYMMETRIC SIMPLE EXCLUSION: THE NAVIER-STOKES CORRECTION

R. Esposito[1], R. Marra[2] and H.T. Yau[3]

1. Dipartimento di Matematica, Università di Roma Tor Vergata, Via della Ricerca Scientifica 00133 Roma, Italy.
2. Dipartimento di Fisica, Università di Roma Tor Vergata, Via della Ricerca Scientifica 00133 Roma, Italy.
3. Courant Institute of Mathematical Sciences, New York University, NY, NY, 10012.

PROBLEM AND RESULTS.

We consider the simple exclusion process (SEP) on a cubic sublattice $\Lambda_L \subset \mathcal{Z}^d$ of size $2L+1$, with periodic boundary conditions. The dynamics is the following: let e denote one of the $2d$ possible directions in $\mathcal{Z}^d$. A particle in the site x, independently of the others, waits for an exponential time and jumps, with probability proportional to $p_e \geq 0$, to the site $x+e$, if it is empty; otherwise stays in x and the process starts again. We denote by $\eta_x(\tau) = 0, 1$ the number of particles in the site x at time τ and by $\mathcal{L}$ the generator of the process: $\mathcal{L}f = \sum_b \mathcal{L}_b f$, the sum running on the set of all oriented bonds $b = (x, y)$ in $\mathcal{Z}^d$ such that $y - x = e$ and

$$\mathcal{L}_b f = p_e \eta_x [f(\eta^b) - f(\eta)] \tag{1}$$

with

$$(\eta^b)_z = (\eta^{x,y})_z = \begin{cases} \eta_y, & z = x \\ \eta_x, & z = y \\ \eta_z, & \text{otherwise.} \end{cases}$$

It is convenient to choose the normalization $p_{-e} + p_e = 2$ for all e.

In the case $p_e = p_{-e}$ for all e (symmetric simple exclusion process) the hydrodynamical limit is non trivial only in the diffusive scaling, $z = \varepsilon x$, $t = \varepsilon^2 \tau$, where x and τ are the microscopic space and time, while z and t are the macroscopic ones, and $\varepsilon = L^{-1}$. Under this scaling, as $\varepsilon \to 0$ the limiting equation reduces to the diffusion equation with diffusion coefficient 1. The case $p_e \neq p_{-e}$, called asymmetric simple exclusion process (ASEP), is much more difficult.

Define the product measure in Λ_L, with chemical potential β, as

$$d\mu_{\beta,L} = Z_\beta^{-1} \exp[\beta \sum_{x \in \Lambda_L} \eta_x], \quad \eta_x \in \{0, 1\}, \tag{2}$$

Z_β being the normalization. $\mu_{\beta,L}$ is invariant for the dynamics because of the periodic b.c., but the dynamics is not reversible with respect to it. The behavior of the system is non trivial already in the hyperbolic (Euler) scaling ($\tau = \varepsilon^{-1} t$) and is given by the non viscous Burgers equation (see ref.[1])

$$\partial_t \rho + \delta \cdot \nabla_z [\rho(1-\rho)] = 0, \tag{3}$$

$\delta_e = p_e - p_{-e}$. Since the system moves already on the Euler scale, it is not clear the meaning of the diffusive limit for this system. For weak asymmetries, namely for the case δ_e of order ε, in the diffusive scaling the process converges to a diffusion with a non linear drift [2,3]. However this case is a small perturbation of the symmetric one and does not manifest the asymmetric nature of the problem. In fact the diffusive constant so obtained is still 1.

We consider[4] the ASEP with δ of order 1, but in a restricted class of initial data for which there is a different mechanism to get a well defined diffusive limiting behavior. This is suggested by the analogy with the study of the incompressible regime for a fluid[5]. Suppose that the initial density $\rho_0(\varepsilon x)$ is of the form

$$\rho_0(\varepsilon x) = \theta - \varepsilon u_0(\varepsilon x)$$

with $\theta \in (0,1)$. Then, assuming that this form of the density persists at time t, the macroscopic current $J = \delta[\rho(1-\rho)]$ becomes

$$J = \delta\theta(1-\theta) - \varepsilon v u - \varepsilon^2 \delta u^2, \quad v = (1-2\theta)\delta.$$

The equation for the field u is expected to be

$$\partial_t u + \varepsilon^{-1} v \cdot \nabla_z u + \delta \cdot \nabla_z u^2 = \sum_{i,j=1}^{d} D_{i,j} \frac{\partial^2 u}{\partial z_i \partial z_j} \tag{4}$$

with $D_{i,j}$ some diffusion matrix. A change of reference frame

$$m(z,t) = u(z + \varepsilon^{-1} v t, t)$$

allows to remove the diverging term and the equation for $m(z,t)$ is the viscous Burgers equation:

$$\partial_t m + \delta \cdot \nabla_z m^2 = \sum_{i,j=1}^{d} D_{i,j} \frac{\partial^2 m}{\partial z_i \partial z_j} \tag{5}$$

This suggests us to assume an initial measure with density w.r.t $\mu_{beta,L}$ given by

$$\psi_0 = Z_\lambda^{-1} \exp\{\varepsilon \sum_x \lambda(\varepsilon x) \eta_x\} \tag{6}$$

for some smooth function $\lambda(z)$, Z_λ being the normalization. Since the product measure with constant chemical potential is invariant, it is natural to think of $\psi_0 d\mu_{\beta,L}$ as a *local equilibrium* for the system.

The average occupation number in the initial local equilibrium state is, up to terms of higher order in ε, $\theta - \varepsilon m_0(\varepsilon x)$, with $\theta = e^\beta/(1+e^\beta)$ and $m(z) = -\theta(1-\theta)\lambda(z)$. By above heuristic arguments one expects that, starting with this initial datum, the occupation number at time t, in the moving frame, differs from θ by $\varepsilon m(z,t)$ with $m(z,t)$ solving (5) with initial datum $m_0(z)$. To state our result we introduce the empirical measure

$$\nu_\epsilon(z,t) = \epsilon^{d-1} \sum_x \delta(z - \epsilon x)(\theta - \eta_x(\varepsilon^{-2} t)).$$

and $\hat\nu_\epsilon(z,t) = \nu_\epsilon(z - \epsilon^{-1} v t, t)$. Next theorem states the weak convergence of $\hat\nu_\epsilon(z,t)$ to $m(z,t)$, if $d \geq 3$.

THEOREM 1.1 *For $d \geq 3$, $\hat\nu_\varepsilon(z,t)$ converges weakly in probability to $m(z,t)$ solving the equation (5). The diffusion matrix satisfies $1 < D < \infty$ (as a matrix) and it is formally given by the Green-Kubo formula.*